


## WEST Search History

DATE: Thursday, October 23, 2003

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=USPT,PGPB,JPAB,EPAB,DWPI; THES=ASSIGNEE; PLUR=YES; OP=ADJ</i>			
L4	HCV 3.0 and L3	1	L4
L3	L2 and HCV adj ELISA	32	L3
L2	Ortho	64675	L2
L1	ELISA3	0	L1

END OF SEARCH HISTORY



Entrez	PubMed	Nucleotide	Protein	Genome	Structure	PMC	Journals	B
Search	PubMed	<input checked="" type="checkbox"/> for microplate sensitized with an antigen					Preview	Go
		Limits	Preview/Index	History	Clipboard	Details		

- Search History will be lost after eight hours of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.

Entrez  
PubMed

Search	Most Recent Queries	Time	Result
#9	Search <b>microplate sensitized with an antigen</b>	12:25:08	<u>7</u>
#8	Search <b>microplate sensitized</b>	12:25:00	<u>17</u>
#1	Search <b>plate sensitized</b>	12:24:17	<u>53</u>


PubMed  
Services

Clear History

Related  
Resources

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
Department of Health & Human Services  
[Freedom of Information Act](#) | [Disclaimer](#)

Oct 24, 2003 09:54 AM



Entrez	PubMed	Nucleotide	Protein	Genome	Structure	PMC	Journals	Books
Search	PubMed	<input checked="" type="checkbox"/> for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details		

Display	Abstract	<input checked="" type="checkbox"/>	Show: 20	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Send to	Text	<input type="checkbox"/>
---------	----------	-------------------------------------	----------	--------------------------	------	--------------------------	---------	------	--------------------------

☐ 1: J Clin Pathol. 1976 Feb;29(2):150-3.

[Related Articles, Links](#)

Entrez  
PubMed

## A microplate enzyme-immunoassay for toxoplasma antibody.

Voller A, Bidwell DE, Bartlett A, Fleck DG, Perkins M, Oladehin B.

PubMed  
Services

A new test for the detection and measurement of toxoplasma antibody is described. Test sera are reacted with **antigen-sensitized wells** in micro-haemagglutination plates. Any attached antibody is shown by the addition of an enzyme-labelled antiglobulin followed by assay of the enzyme reaction with its substrate. The test is easy to carry out on a large scale, and there is a positive correlation between the results and dye test and haemagglutination test titres.

PMID: 932215 [PubMed - indexed for MEDLINE]

Related  
Resources

Display	Abstract	<input checked="" type="checkbox"/>	Show: 20	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Send to	Text	<input type="checkbox"/>
---------	----------	-------------------------------------	----------	--------------------------	------	--------------------------	---------	------	--------------------------

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)



Entrez	PubMed	Nucleotide	Protein	Genome	Structure	PMC	Journals	Books	
Search	PubMed	for						Go	Clear
Limits	Preview/Index	History	Clipboard	Details					

Display	Abstract	Show: 20	Sort	Send to	Text
---------	----------	----------	------	---------	------

☐ 1: J Immunol Methods. 1988 Oct 4;113(1):17-24.

[Related Articles, Links](#)

Entrez  
PubMed

## A comparison of immunological methods for the detection of *Trichinella spiralis* antigen.

Choy WF, Lim PL, Ng MH.

PubMed  
Services

Department of Microbiology, University of Hong Kong.

Related  
Resources

Eight immunological methods all using the same monoclonal antibody reagent were compared for the detection of *Trichinella spiralis* antigen. These were based on: (1) the direct adsorption of the antigen to the immunoadsorbent (nitrocellulose membrane, polyvinyl chloride strip or microplate); (2) capture of the antigen by **antibodies pre-sensitized on the immunoadsorbent**; and (3) latex agglutination. The methods found suitable were: (a) capture radioimmunoassay (capture-RIA) (sensitivity: less than 0.5 microgram/ml antigen); (b) direct enzyme immunoassay (direct-ELISA) (less than 0.5 microgram/ml); (c) tube latex agglutination test (2.2 micrograms/ml); and (d) direct immunodot assay (8.8 micrograms/ml). However, the performance of the direct-ELISA was greatly affected by the presence of each of three extraneous substances (bovine serum albumin (BSA), lipopolysaccharide (LPS), normal swine muscle homogenate (NSM) added to the antigen sample. The direct immunodot assay was also affected by the presence of BSA or LPS, whereas both the capture-RIA and the tube latex agglutination methods were affected by the presence of NSM only. The findings are discussed with a view of detecting *T. spiralis* larvae directly from pork samples by immunological means.

PMID: 3049823 [PubMed - indexed for MEDLINE]

Display	Abstract	Show: 20	Sort	Send to	Text
---------	----------	----------	------	---------	------

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)

Oct 20 2003 06:54:00